

COLYTE 3 DAY BOWEL PREP

The following instructions are your physician's specific instructions. Please follow the instructions carefully to ensure a successful prep.



You can reach Your Patient Advisor with non-medical prep questions at: **800.349.0285**

You can reach your physician's office at: **626.359.3330 (ARCADIA) 626.963.2490 (GLENORA)**

Special Notes: Fill prescription for one 4L preparation. Do **NOT** mix container with water yet.

Buy 8 Generic Bisacodyl tablets (5mg each tablet). This may be dispensed with your 4L preparation, however if it is not, there are several brands of Bisacodyl available. If you are unsure, ask the pharmacist for assistance.

SEE PAGE 5 FOR DETAILED MEDICATION INSTRUCTIONS.

7 days prior to procedure	3 days prior to procedure	2 days prior to procedure	1 day prior to procedure	Day of procedure
<p>Review your prep instructions thoroughly.</p> <p>STOP: iron pills because it leaves a dark residue inside your colon.</p> <p>STOP: ALL vitamins and supplements because they can leave residues and debris inside you.</p> <p>STOP: fiber supplements (Metamucil, Benefiber, Citrucel, Psyllium, etc.)</p>	 <p>STOP eating any raw vegetables or vegetables containing seeds, corn, popcorn, nuts, and seeds and stop any fiber supplements until after the procedure.</p> <p>Purchase clear liquid diet items, ointment, reading material, etc.</p> <p>Prep at a Glance 5:00 pm take 2 Bisacodyl</p>	 <p>Continue a low fiber diet. Remember to avoid foods with nuts and seeds.</p> <p>Confirm arrangements with your driver.</p> <p>Prep at a Glance 5:00 pm take 2 Bisacodyl</p>	<p>NO SOLID FOODS, CLEAR LIQUIDS* ONLY upon rising, until after your procedure. A clear liquid diet is necessary for a colonoscopy.</p> <p>Prep at a Glance 3:00 pm take 2 Bisacodyl 6:00 pm take 96oz of prep solution</p> <p>Drink plenty of water and liquids throughout the day to avoid dehydration.</p> <p>Detailed, step-by-step instructions continue on page 3.</p>	<p>4 hours prior to your arrival time Drink remaining 32oz of prep solution.</p> <p>Nothing by mouth 3 hours prior to your procedure.</p> <p>NO GUM OR HARD CANDY</p> <p>YOU MAY BRUSH YOUR TEETH BUT DO NOT SWALLOW</p> <p>*If you take daily medication, you may take it with SMALL SIPS OF WATER ONLY, at least 3 hours before your procedure.</p>

***Clear Liquid Diet Details: NO RED or PURPLE. NO DAIRY**

Approved

- Sodas, coffee, tea
- Clear juices, fitness waters
- Popsicles without pulp
- Chicken, vegetable and beef broth
- Gelatin

Avoid

- No milk/dairy
- No juices with pulp
- NO RED or PURPLE**



Bowel Prep Frequently Asked Questions

What is a clear stool?

A clear stool can have a slight tint of yellow or brown. It will be completely transparent, and will not contain any solid matter.

I am not having bowel movements, what should I do?

Bowel movements can take up to 5-6 hours after beginning the prep to start. Be patient, continue to drink liquids. If you have not had a bowel movement by midnight the night prior to your procedure, you will need to reach the on call physician for further instructions.

The prep is making me nauseous, what should I do?

If you develop nausea or vomiting, slow down the rate at which you drink the solution. Please attempt to drink all of the laxative solution even if it takes you longer. If vomiting persists, or you are not able to finish the preparation, stop the preparation and call your physician's office for further instructions.

What are some high fiber foods I should avoid?

Raw fruits and vegetables are typically high in fiber, as well as nuts, seeds, whole grain breads and beans and lentils.

What are some good options for low fiber foods?

Choose white bread and white rice for lower fiber options, as well as pastas made with white flour. Chicken, fish, dairy and eggs are also low in fiber and good choices for foods 2-3 days before you begin your prep.

If I eat popcorn or seeds 3 days before my procedure do I need to reschedule?

You will not need to reschedule your procedure, however the seeds or nuts may cause a difficulty in screening and require a need for rescreening. If you have eaten a large amount of seeds or nuts, you may want to contact the nurse or physician.

Can I drink ALCOHOL on the liquid diet?

Alcohol is not allowed as part of the liquid diet.

Can I continue to be on the liquid diet after I begin consuming the laxatives?

Yes, you may continue the liquid diet until you are directed to discontinue anything by mouth, which is typically 4-6 hours prior to the procedure.

Why do I have to wake up at so early for the 2nd dose, can't I take it all the night before?

A split prep has proven to be the most effective for a successful colonoscopy. It is essential that you follow the directions provided with your prep medications.

If I weigh under 100 pounds do I need to take all of the prep?

The liquid amount is not weight dependent. It is important to finish the prep for a successful colonoscopy.

COLYTE 3 DAY BOWEL PREP

Prep Day: Three days before your procedure

1

5:00 PM three days prior to your procedure

Take 2 (5mg) Bisacodyl tablets with a glass of water.

***If no results in 2 hours, please take 2 additional tablets.**



Prep Day: Two days before your procedure

2

5:00 PM two days prior to your procedure

Take 2 (5mg) Bisacodyl tablets with a glass of water.

***If no results in 2 hours, please take 2 additional tablets.**



Prep Day: The day before your procedure

No solid foods upon rising, until after your procedure. Drink plenty of water throughout the day to avoid dehydration.

IF YOU ARE DIABETIC, PLEASE REVIEW THE FOLLOWING INSTRUCTIONS:

- If you take pills or other oral diabetic medication, stop your medication the day before and the morning of the procedure.
- If you take insulin or other injectable diabetic medications, take **1/3** of the usual dose at the normal time. If glucose is over 250, you can take 5 units of regular or 70/30 insulin (but not Lispro/Short Acting Insulin).
- If you have a short acting insulin or sliding scale insulin, please continue course according to blood sugar level.
- If your blood sugar goes below 80 or you have symptoms of hypoglycemia, please drink apple juice.
- **If you get your insulin through an implanted pump, please check with your diabetes prescribing physician.**
- Hold BYETTA and RICTOZA.

**PLEASE NOTE YOU CAN ALSO START THE PROCESS BELOW WHEN YOU GET HOME FROM WORK.
PLEASE ADJUST THE TIMES ACCORDINGLY.**

Mixing Your Bowel Preparation:

- Colyte/TriLyte/Nulytely/Golytely are common names for PEG 3350 solution.
- **MIX** the preparation using these instructions, **NOT** the instructions on the bottle.

3

Before you go to work the day prior to your procedure

Add 2 inches of lukewarm water to the bottom of the 4L preparation container, mix well then add cold water to the "fill line" and chill in the refrigerator.

DO NOT USE THE INSTRUCTIONS FOR MIXING ON THE BOTTLE



COLYTE 3 DAY BOWEL PREP

4

3:00 PM the day prior to your procedure
Take 2 (5mg) Bisacodyl tablets with a glass of water.



5

6:00 PM the evening prior to your procedure
Begin drinking the preparation. Drink an 8 oz. glass of the solution every 20-30 minutes **until 3/4 the container is gone** (total of **12** glasses).



Prep Day: The day of your procedure

6

4 hours prior to your arrival time the morning of your procedure

Drink an 8 oz. glass of the solution every 20-30 minutes **until the remainder of the container is gone** (total of **4** glasses)

Shake the container before pouring each dose.



THE DAY OF YOUR PROCEDURE

- **NOTHING BY MOUTH 3 HOURS PRIOR TO THE PROCEDURE, WITH THE EXCEPTION OF YOUR PREPARATION.**
- **NOTHING TO EAT OR DRINK UNTIL AFTER YOUR PROCEDURE.**
- Take your usual medications that you normally take the morning of the procedure with a small sip of water, except the one you have been instructed to hold.

ADDITIONAL IMPORTANT INFORMATION

- **IT IS ESSENTIAL THAT A RESPONSIBLE ADULT DRIVES YOU HOME FROM THE EXAM. YOU CANNOT RIDE HOME ON A BUS, TAKE A TAXI, OR WALK HOME BY YOURSELF.**
- Bring with you your insurance card and the pamphlet about your rights and responsibilities as a patient to your procedure.
- Wear comfortable clothing and socks to keep your feet warm.
- Do **not** wear jewelry, tight belts, pantyhose, cologne or perfume.
- Leave money and other valuables at home.
- Your judgment and reflexes may be impaired due to the sedatives you will receive. You will not be able to drive, operate machinery, sign legal documents, drink alcoholic beverages, or go to work until the following day.

COLYTE 3 DAY BOWEL PREP

- If you take prescription blood pressure, heart, lung, seizure, (narcotic) pain or psychiatric medication daily, please take your daily dose(s) as prescribed with a sip of water, including your procedure day.

If you have any questions or concerns about the preparation, please contact Your Patient Advisor by calling 800-349-0285 or emailing support@yourpatientadvisor.com.

If you have any medical questions or concerns, please contact one of our offices at 626-359-3330 (Arcadia) or 626-963-2490 (Glendora).

MEDICATION RESTRICTION INFORMATION

SPECIAL NOTE: Please make sure you have obtained permission from your cardiologist or internist to hold these drugs. If you have a coronary stent or other important condition, stopping these drugs could lead to serious consequences.

- If you take **aspirin** on a daily basis, for medical reasons, please continue taking it.
- If you take **Plavix**, please continue taking it.*
- If you take **Coumadin** or **Warfarin**, please hold for **3 days** prior to your procedure.*
- If you take **Pradaxa**, **Xarelto** or **Eliquis**, please hold for **2 days** prior to your procedure.*
- If you take any other blood thinner, please make sure you get specific dosing instructions from your prescribing physician.
- If your blood thinner anticoagulation prescription is managed by a Coumadin clinic, please inform them of planned procedure and ask for advice.
- Tylenol (acetaminophen) is fine to use, as needed.